arXiv:1609.02348v1 [math.AG] 8 Sep 2016

AUTOMORPHISMS OF SALEM DEGREE 22 ON SUPERSINGULAR K3 SURFACES OF HIGHER ARTIN INVARIANT - A SHORT NOTE

SIMON BRANDHORST

Abstract. We give a short proof that every supersingular K3 surface in odd characteristic has an automorphism of Salem degree 22. The proof relies on the case $\sigma = 1$ and the cone theorem.

1. Introduction

A Salem number is a real algebraic integer $\lambda > 1$ which is conjugate to $1/\lambda$ and all whose other conjugates lie on the unit circle. We call its minimal polynomial a Salem polynomial. If X is a projective K3 surface over an algebraically closed field k and $f\colon X \to X$ an automorphism, then the characteristic polynomial $\chi(f^*|H^2_{\acute{e}t}(X, \mathbb{Q}_l))$ ($l \neq \operatorname{char} k$) factors as a product of cyclotomic polynomials and at most one Salem polynomial [4]. We call the degree of the Salem factor $s(x)$ the Salem degree of f. It is well known that $\ker s(f^*|H^2_{\acute{e}t}(X, \mathbb{Q}_l))$ is contained in the Néron-Severi group $\mathrm{NS}(X) \otimes \mathbb{Q}_l$ of X. In particular we can bound the Salem degree of an automorphism by the Picard number $\rho(X)$, which is at most $h^{1,1}(X) = 20$ in characteristic 0. However in positive characteristic $\rho(X) = 22$ is possible. These are the so called supersingular K3 surfaces. This observation leads to the interesting feature that an automorphism of Salem degree 22 is not geometrically liftable to characteristic zero (see [3] for details). The existence of such automorphisms in any characteristic is established in the following

Theorem 1.1. [1, 2, 3, 5, 9, 10] *The supersingular K3 surface X/k, $k = \overline{k}$, $\operatorname{char} k > 0$, of Artin invariant one has an automorphism of Salem degree 22.*

Supersingular K3 surfaces are classified by their Artin invariant $1 \leq \sigma \leq 10$. For fixed Artin invariant σ they form a family of dimension $\sigma - 1$. A recent preprint generalizes Theorem 1.1 above to K3 surfaces in odd characteristic of *any* Artin invariant.

Theorem 1.2. [10, $p = 3$][11, $p > 3$] *Every supersingular K3 surface Y/k over an algebraically closed field $k = \overline{k}$ of characteristic $\operatorname{char} k = p > 2$ has an automorphism of Salem degree 22.*

The proof for $p > 3$ involves genus one fibrations. We take a different point of view and rely instead on the $\sigma = 1$ case and the Cone Theorem 2.1. In characteristic 2 the cone theorem is proved provided that $\sigma(Y) < 10$. Hence we get an extension of the above theorem to this case as well.

Acknoledgements. I thank Hélène Esnault, Víctor González-Alonso, Keiji Oguiso, Matthias Schütt, and Xun Yu for discussions and comments on this work.

2. The proof

Theorem 2.1 (Cone Theorem)**.** [6, Thm. 6.1] *Let X be a K3 surface over an algebraically closed field of characteristic $p > 3$. Let $\Gamma(X) \subseteq O(\mathrm{NS}(X))$ be the subgroup preserving the nef cone. Then the natural map $\mathrm{Aut}(X) \to \Gamma(X)$ has finite kernel and cokernel.*

Over $\mathbb{C}$ this is well known. Then for K3 surfaces of finite height in arbitrary characteristic the proof works through lifting to characteristic zero. For supersingular K3 surfaces it uses the crystalline Torelli Theorem, which is proved for $p > 3$ or $p = 2$ and $\sigma < 10$ or $p = 3$ and $\sigma < 6$ (see [7] and the end of [8]).
Since any power of a Salem number of degree d remains a Salem number of this degree, passing to a finite index subgroup does not change the maximal occurring Salem degree of a matrix group. Combining this with the Cone Theorem, we get that the maximum occurring Salem degree of an automorphism of X depends only on $\Gamma(X)$. Now $\Gamma(X)$ depends up to conjugation by an element of the Weyl group only on the isometry class of $\mathrm{NS}(X)$. In particular the maximal Salem degree of an automorphism of X depends only on $\mathrm{NS}(X)$.

Lemma 2.2. *Let $N \subseteq L$ be two lattices of the same rank and $G \subseteq O(L)$ a subgroup. Then*

$$[G : O(N) \cap G] < \infty$$

where we view $O(N)$ and $O(L)$ as subgroups of $O(N \otimes \mathbb{R})$.

Proof. Since the ranks coincide, the index $n = [L : N]$ is finite and

$$nL \subseteq N \subseteq L.$$

Any isometry of L preserves nL hence we get a map

$$\varphi\colon G \to \mathrm{Aut}(L/nL).$$

Set $K = \ker \varphi$, which is a finite index subgroup of G. To see that $K \subseteq O(N)$ as well, recall that an isometry f of $O(nL)$ extends to $O(N)$ iff $f(N/nL) = N/nL$. Indeed, $f|L/nL = id|L/nL$ for $f \in K$, by definition. □

Theorem 2.3. *Let X, Y be two K3 surfaces over an algebraically closed field k,* $\mathrm{char}\, k > 3$. *Suppose that $\rho(X) = \rho(Y)$ and there is an isometric embedding*

$$\iota : \mathrm{NS}(Y) \hookrightarrow \mathrm{NS}(X).$$

Then $\mathrm{sdeg}(X) \leq \mathrm{sdeg}(Y)$ *where* $\mathrm{sdeg}(X) = \max\{$*Salem degree of* $f | f \in \mathrm{Aut}(X)\}$.

Proof. Denote by $\mathrm{Nef}(X)$ and $\mathrm{Nef}(Y)$ the nef cones of X and Y. Any chamber of the positive cone of $\mathrm{NS}(X)$ is contained in the image of a unique chamber of the positive cone of $\mathrm{NS}(Y)$. Since the Weyl group acts transitively on the chambers, we can find an element $r \in W(\mathrm{NS}(X))$ of the Weyl group such that $\mathrm{Nef}(X) \subset \iota'_{\mathbb{R}}(\mathrm{Nef}(Y))$ where $\iota' = r \circ \iota$. To ease notation we identify $\mathrm{NS}(Y)$ with its image under ι'. By the preceding Lemma $[\Gamma(X) : \Gamma(X) \cap O(\mathrm{NS}(Y))]$ is finite, and since $\mathrm{Nef}(X) \subseteq \mathrm{Nef}(Y)$, we get that $\Gamma(X) \cap O(\mathrm{NS}(Y)) \subseteq \Gamma(Y)$. Now, by the cone Theorem 2.1 and the discussion after it we can conclude with

$$\mathrm{sdeg}(X) = \mathrm{sdeg}(\Gamma(X)) = \mathrm{sdeg}(\Gamma(X) \cap O(NS(Y))) \leq \mathrm{sdeg}(\Gamma(Y)) = \mathrm{sdeg}(Y).$$

□

Proof of Theorem. 1.2. It is well known that if X/k and Y/k are supersingular K3 surfaces with $\sigma(X) \leq \sigma(Y)$, then $\mathrm{NS}(Y) \hookrightarrow \mathrm{NS}(X)$. Combining the $\sigma = 1$ case and the previous theorem we get that $22 = \mathrm{sdeg}(X) \leq \mathrm{sdeg}(Y) \leq 22$. □

The converse inequality in Theorem 2.3 is false in general. This was pointed out to me by K. Oguiso and X. Yu.

References

[1] J. Blanc and S. Cantat, *Dynamical degrees of birational transformations of projective surfaces*, *J. Amer. Math. Soc.* **29** (2016) 415–471.

[2] S. Brandhorst, *Dynamics on supersingular K3 surfaces and automorphisms of Salem degree 22*, *Nagoya Math. J., to appear* , [arXiv:1507.02092].

[3] H. Esnault and K. Oguiso, *Non-liftability of automorphism groups of a K3 surface in positive characteristic*, *Math. Ann.* **363** (2015) 1187–1206.

[4] H. Esnault and V. Srinivas, *Algebraic versus topological entropy for surfaces over finite fields*, *Osaka J. Math.* **50** (2013) 827–846.

[5] H. Esnault, K. Oguiso and X. Yu, *Automorphisms of elliptic K3 surfaces and Salem numbers of maximal degree*, *J. Alg. Geom., to appear* , [arXiv:1411.0769].

[6] M. Lieblich and D. Maulik, *A note on the cone conjecture for k3 surfaces in positive characteristic*, arXiv:1102.3377v3.

[7] A. Ogus, *A crystalline Torelli theorem for supersingular K3 surfaces*, in *Arithmetic and geometry, Vol. II*, vol. 36 of *Progr. Math.*, pp. 361–394. Birkhäuser Boston, Boston, MA, 1983.

[8] A. N. Rudakov and I. R. Shafarevich, *Surfaces of type K3 over fields of finite characteristic*, in *Current problems in mathematics, Vol. 18*, pp. 115–207. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Informatsii, Moscow, 1981.

[9] M. Schütt, *Dynamics on supersingular K3 surfaces*, *Comm. Math. Helvetici, to appear* , [arXiv:1502.06923].

[10] I. Shimada, *Automorphisms of supersingular K3 surfaces and Salem polynomials*, *Exp. Math.* **25** (2016) 389–398.

[11] X. Yu, *Elliptic fibrations on K3 surfaces and Salem numbers of maximal degree*, arXiv:1605.09260.

Insitut für Algebraische Geometrie, Leibniz Universität Hannover, Welfengarten 1, 30167 Hannover, Germany

E-mail address: brandhorst@math.uni-hannover.de